Author Index

Abe, T., see Ogawa, K. 188
Abián, J., see Cárdenas, A. 83
Abumrad, N.N., see Brown, L.L. 370
Agarwal, G.S., see Ramachandran, P.K. 239
Aguilar, M.I.
—, Hart, S.J. and Calder, I.C.
Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair high-performance liquid chromatography 315
Akbari, A.
—, Gordon, B.J., Bush, P.B. and Moore, J.N.
Determination of xylazine in blood components using high-performance liquid chromatography 207
Amantea, M.A.
— and Narang, P.K.
Improved procedure for quantitation of omeprazole and metabolites using reversed-phase high-performance liquid chromatography 216
Antonucci, A., see Pecci, L. 183
Baba, S., see Sasaki, Y. 93
Bajeva, S.K., see Padmalatha Devi, K. 229
Baldessarini, R.J., see Lampen, P. 283
Bayne, W.F., see Matuszewski, B.K. 295
Beaune, P.H.
—, Cresteil, T., Flinois, J.-P. and Leroux, J.-P.
Hydrophobic chromatography: a one-step method for the purification of human liver microsomal epoxide hydrolase 169
Becker, T.A., see Brown, L.L. 370
Belsner, K., see Weber, C. 345
Benet, L.Z., see Lee, F.W. 259
Bernstein, J.R., see Miller, B.J. 438
Bhargava, V.O.
—, Emodi, S. and Hirate, J.
Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 212
Birkett, D.J., see Chan, K. 103
Bizzi, B., see De Cristofaro, R. 376
Bleau, G., see Vézina, D. 385
Brashear, W.T., see Kuhnert, B.R. 392
Brendel, E.
—, Meineke, I., Henne, E.-M., Zschunke, M. and De Mey, C.
Sensitive high-performance liquid chromatographic determination of pseudoephedrine in plasma and urine 406
Brown, H., see Holland, O.B. 148
Brown, L.L.
—, Williams, P.E., Becker, T.A., Ensley, R.J., May, M.E. and Abumrad, N.N.
Rapid high-performance liquid chromatographic method to measure plasma leucine: importance in the study of leucine kinetics in vivo 370
Buckley, D.G., see Whelpton, R. 223
Bulbena, O., see Cárdenas, A. 83
Burke, J.T., see Krstulović, A.M. 417
Bush, P.B., see Akbari, A. 207
Calder, I.C., see Aguilar, M.I. 315
Cárdenas, A.
—, Abián, J., Bulbena, O., Roselló, J. and Gelpi, E.
Determination of oxidation products of N-phenyllinoleamide: Spanish toxic oil syndrome studies 83
Castagnola, M., see De Cristofaro, R. 376
Cavallini, D., see Pecci, L. 183
Chan, K.
—, Miners, J.O. and Birkett, D.J.
Direct and simultaneous high-performance liquid chromatographic assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 103
Cheymol, G., see Poirier, J.-M. 431
Constanzer, M.L., see Matuszewski, B.K. 295
Costa, M., see Pecci, L. 183
Cresteil, T., see Beaune, P.H. 169
Cristofaro, R. De, see De Cristofaro, R. 376
Davis, L.J., see Hoskins, J.A. 155
D'Costa, M., see Diamandis, E.P. 25
Debackere, M., see Delbeke, F.T. 194
De Cristofaro, R.
—, Landolfi, R., Bizzi, B. and Castagnola, M.
Human platelet glycocalicin purification by phenyl boronate affinity chromatography coupled to anion-exchange high-performance liquid chromatography 376
Delbeke, F.T.
—, Debackere, M., Desmet, N. and Maertens, F.
Qualitative gas chromatographic and gas chromatographic–mass spectrometric screening for β-blockers in urine after solid-phase extraction using Extrelut-1 columns 194
De Mey, C., see Brendel, E. 406
Desmet, N., see Delbeke, F.T. 194

Devi, K. Padmalatha, see Padmalatha Devi, K. 229
Deyl, Z.
—, Horáková, M. and Macek, K.
Column liquid chromatographic separation of collagen α-chain polymers on sepiolite 162
Diamandis, E.P.
— and D'Costa, M.
Selective determination of urinary free cortisol by liquid chromatography after solid-state extraction 25
Dube, P., see Holland, O.B. 148
Echizen, H.
—, Itoh, R. and Ishizaki, T.
Microassay of free 3-methoxy-4-hydroxyphenylglycol in plasma using high-performance liquid chromatography with electrochemical detection 351
Emodi, S., see Bhargava, V.O. 212
Endoh, Y.S.
—, Yoshimura, H. and Sasaki, N.
High-performance liquid chromatographic determination of bromofenofos and its metabolite in rat plasma 202
Ensley, R.J., see Brown, L.L. 370
Fasolato, C.
— and Galzigna, L.
Assay of γ-L-glutamylcyclotransferase activity in rat brain synaptosomes by high-performance liquid chromatography 381
Fathi, M., see Padmalatha Devi, K. 229
Fayolle, C.
— and Fillion, G.
Reversed-phase liquid chromatographic determination of cyclic adenosine 3′,5′-monophosphate in rat brain cortex 177
Fillion, G., see Fayolle, C. 177
Flamini, G.
—, Ria, F., Pizzigallo, E. and Gignina, S.
Chromatographic and electrophoretic studies of immune complexes in non-A, non-B hepatitis 55
Flinois, J.-P., see Beaune, P.H. 169
Franklin, R.B., see Miller, B.J. 438
Fujiwara, Y., see Ogawa, K. 188
Galzigna, L., see Fasolato, C. 381
Gandhe, B.R., see Ramachandran, P.K. 239
Gelpi, E., see Cárdenas, A. 83
Gianviti, J.M., see Krstulović, A.M. 417
Gietl, Y.
—, Spahn, H. and Mutschler, E.
Simultaneous determination of *R*- and *S*-prenylamine in plasma and urine by reversed-phase high-performance liquid chromatography 305
Gignina, S., see Flamini, G. 55
Goldstein, D.S., see Hovevey-Sion, D. 141
Gomez-Sanchez, C., see Holland, O.B. 148
Gopalan, N., see Ramachandran, P.K. 239
Gordon, B.J., see Akbari, A. 207
Gozel, P., see Roach, M.C. 129
Grizzle, W., see Jenkins, R.L. 249
Harrison, P.M., see Kech, A.C. 234
Hart, S.J., see Aguilar, M.I. 315
Harvey-White, J., see Hovevey-Sion, D. 141
Hatsumura, H., see Sayo, H. 449
Hayashi, N., see Suzuki, Y. 33
Henne, E.-M., see Brendel, E. 406
Hirate, J., see Bhargava, V.O. 212
Hirooka, T., see Naoi, M. 75
Höller, M., see Weber, C. 345
Hoffman, N.E., see Rustum, A.M. 121
Holland, O.B.
—, Brown, H., Risk, M., Paone, R., Dube, P. and Gomez-Sanchez, C.
Simplified method for the determination of plasma 18-hydroxycorticosterone 148
Holthuis, J.J.M., see Van der Horst, F.A.L. 267
Horáková, M., see Deyl, Z. 162
Horst, F.A.L. van der, see Van der Horst, F.A.L. 267
Hoskins, J.A.
— and Davis, L.J.
Analysis of the isomeric tyrosines in mammalian and avian systems using high-performance liquid chromatography with fluorescence detection 155
Hosokawa, M., see Sayo, H. 449
Hovevey-Sion, D.
—, Harvey-White, J., Kopin, I.J. and Goldstein, D.S.
Measurement of homovanillic acid in small volumes of plasma using liquid chromatography with electrochemical detection 141
Ishii, M.
— and Ogata, H.
Determination of rifampicin and its main metabolites in human plasma by high-performance liquid chromatography 412
Ishizaki, T., seee Echizen, H. 351
Itoh, R., see Echizen, H. 351
Jaillon, P., see Poirier, J.-M. 431
Jenkins, R.L.
—, McDaniel, H.G., Grizzle, W., Parrish, S.W. and McDaniel, H.B.

Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by high-performance liquid chromatography 249
Jonas, G., see Whelpton, R. 223
Kapetanovic, I.M., see Torchin, C.D. 444
Karhu, I.
—, Mahonen, A. and Palvimo, J.
Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase liquid chromatography: protein levels in rat tissues during postnatal development 65
Kaushik, M.P., see Ramachandran, P.K. 239
Kawase, M., see Ohmori, S. 15
Keech, A.C.
—, Harrison, P.M. and McLean, A.J.
Simple extraction of atenolol from urine and its determination by high-performance liquid chromatography 234
Kissmeyer-Nielsen, A.-M.
Determination of 6β-bromopenicillanic acid (brobactam) in human serum and urine by high-performance liquid chromatography 425
Kojima, T., see Naoi, M. 75
Kondoh, M., see Naoi, M. 75
Kopin, I.J., see Hovevey-Sion, D. 141
Krstulović, A.M.
—, Gianviti, J.M., Burke, J.T. and Mompon, B.
Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chiral β-cyclodextrin stationary phase 417
Kuhnert, B.R.
—, Brashear, W.T. and Syracuse, C.D.
Measurement of alphaprodine by selected-ion monitoring 392
Kupferberg, H.J., see Torchin, C.D. 444
Lam, H.R.
— and Tarding, F.
Specific high-performance liquid chromatographic method for the determination of creatinine in rat plasma 358
Lampen, P.
—, Neumeyer, J.L. and Baldessarini, R.J.
High-performance liquid chromatographic separation and electrochemical or spectrophotometric determination of *R*(−)N-*n*-propylnorapomorphine and *R*(−)10,11-methylenedioxy-N-*n*-propylnoraporphine in primate plasma 283
Landolfi, R., see De Cristofaro, R. 376
Lee, F.W.
—, Watari, N., Rigod, J. and Benet, L.Z.
Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary gas chromatography with electron-capture detection 259
Leroux, J.-P., see Beaune, P.H. 169
Lohkamp, R., see Weber, C. 345
McDaniel, H.B., see Jenkins, R.L. 249
McDaniel, H.G., see Jenkins, R.L. 249
Macek, K., see Deyl, Z. 162
McLean, A.J., see Keech, A.C. 234
Maertens, F., see Delbeke, F.T. 194
Mahonen, A., see Karhu, I. 65
Matuszewski, B.K.
—, Constanzer, M.L. and Bayne, W.F.
Determination of 2,3-dihydro-6-[3-(2-hydroxymethyl)phenyl-2-propenyl]-5-benzofuranol in plasma using liquid chromatography with electrochemical detection 295
May, M.E., see Brown, L.L. 370
Mehta, A.C.
Review. Direct separation of drug enantiomers by high-performance liquid chromatography with chiral stationary phases 1
Meineke, I., see Brendel, E. 406
Mey, C. De, see Brendel, E. 406
Miller, B.J.
—, Tenbarge, J.B., Bernstein, J.R. and Franklin, R.B.
High-performance liquid chromatographic assay of *trans*-(−)-5,5a,6,7,8,9,9a,10-octahydro-6-propylpyrimido[4,5-*g*]quinolin-2-amine dihydrochloride (LY 163502), a potent D-2 agonist, in plasma 438
Miners, J.O., see Chan, K. 103
Mompon, B., see Krstulović, A.M. 417
Moore, J.N., see Akbari, A. 207
Mutoh, T., see Naoi, M. 75
Mutschler, E., see Gietl, Y. 305
Nagatsu, T., see Naoi, M. 75
Nakata, K., see Ohmori, S. 15
Naoi, M.
—, Kondoh, M., Mutoh, T., Takahashi, T., Kojima, T., Hirooka, T. and Nagatsu, T.
Microassay for GM_1 ganglioside β-galactosidase activity using high-performance liquid chromatography 75
Narang, P.K., see Amantea, M.A. 216
Neumeyer, J.L., see Lampen, P. 283
Ogata, H., see Ishii, M. 412
Ogawa, K.
—, Fujiwara, Y., Sugamata, K. and Abe, T.
Thin-layer chromatography of neutral gly-

cosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga_2Cer 188
Ohmori, S.
—, Sumii, I., Toyonaga, Y., Nakata, K. and Kawasa, M.
High-performance liquid chromatographic determination of formate as benzimidazole in biological samples 15
Okayama, A.
Fluorimetric determination of urinary δ-aminolevulinic acid by high-performance liquid chromatography and post-column derivatization 365
Ono, K., see Tanaka, M. 111
Padmalatha Devi, K.
—, Ranga Rao, K.V., Bajeva, S.K., Fathi, M. and Roth, M.
Determination of oxprenolol in plasma by column liquid chromatography 229
Palvimo, J., see Karhu, I. 65
Paone, R., see Holland, O.B. 148
Parrish, S.W., see Jenkins, R.L. 249
Pecci, L.
—, Costa, M., Pinnen, F., Antonucci, A. and Cavellini, D.
Specific method for the determination of 1,4-thiomorpholine-3,5-dicarboxylic acid and its unsaturated analogue lanthionine ketimine 183
Perez, M., see Poirier, J.-M. 431
Pinnen, F., see Pecci, L. 183
Pizzigallo, E., see Flamini, G. 55
Poirier, J.-M.
—, Perez, M., Cheymol, G. and Jaillon, P.
Rapid and sensitive high-performance liquid chromatographic determination of bisoprolol in plasma and urine 431
Ramachandran, P.K.
—, Gandhe, B.R., Venkateswaran, K.S., Kaushik, M.P., Vijayaraghavan, R., Agarwal, G.S., Gopalan, N., Suryanarayana, M.V.S., Shinde, S.K. and Sriramachari, S.
Gas chromatographic studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 239
Ranga Rao, K.V., see Padmalatha Devi, K. 229
Rao, K.V. Ranga, see Padmalatha Devi, K. 229
Ria, F., see Flamini, G. 55
Rigod, J., see Lee, F.W. 259
Risk, M., see Holland, O.B. 148
Roach, M.C.
—, Gozel, P. and Zare, R.N.
Determination of methotrexate and its major metabolite, 7-hydroxymethotrexate, using capillary zone electrophoresis and laser-induced fluorescence detection 129
Rodchenkov, G.M.
—, Uralets, V.P. and Semenov, V.A.
Gas chromatographic and mass spectral study of synthetic corticosteroid metabolism: fluorometholone 399
Romijn, J.C., see Verkoelen, C.F. 41
Roselló, J., see Cárdenas, A. 83
Roth, M., see Padmalatha Devi, K. 229
Rouan, M.C.
Microbore liquid chromatographic determination of cadralazine and cephalexin in plasma with large-volume injection 335
Rustum, A.M.
— and Hoffman, N.E.
Determination of 5-fluorouracil in plasma and whole blood by ion-pair high-performance liquid chromatography 121
Sasaki, N., see Endoh, Y.S. 202
Sasaki, Y.
— and Baba, S.
Simultaneous determination of imipramine desipramine and their deuterium-labelled analogues in biological fluids by capillary gas chromatography–mass spectrometry 93
Sayo, H.
—, Hatsumura, H. and Hosokawa, M.
Simultaneous determination of phenytoin and phenobarbital in human serum by electrochemical immunoassay combined with high-performance liquid chromatography 449
Schalkwijk, W.P. van, see Verkoelen, C.F. 41
Schroeder, F.H., see Verkoelen, C.F. 41
Sekiba, K., see Suzuki, Y. 33
Semenov, V.A., see Rodchenkov, G.M. 399
Shinde, S.K., see Ramachandran, P.K. 239
Spahn, H., see Gietl, Y. 305
Splinter, T.A.W., see Verkoelen, C.F. 41
Sriramachari, S., see Ramachandran, P.K. 239
Sugamata, K., see Ogawa, K. 188
Sumii, I., see Ohmori, S. 15
Suryanarayana, M.V.S., see Ramachandran, P.K. 239
Suzuki, Y.
—, Hayashi, N. and Sekiba, K.
Automated direct assay system for the measurement of sex steroid hormones in serum using high-performance liquid chromatography 33
Syracuse, C.D., see Kuhnert, B.R. 392
Takahashi, T., see Naoi, M. 75

Takegoshi, T., see Tanaka, M. 111
Tanaka, M.
—, Ono, K. and Takegoshi, T.
Determination of the thromboxane synthethase inhibitor 6-(1-imidazolylmethyl)-5,6,7,8,-tetrahydronaphthalene-2-carboxylic acid (DP-1904) in human plasma and urine using solid-phase extraction and high-performance liquid chromatography 111
Tarding, F., see Lam, H.R. 358
Tenbarge, J.B., see Miller, B.J. 438
Torchin, C.D.
—, Kapetanovic, I.M., Yonekawa, W.D. and Kupferberg, H.J.
High-performance liquid chromatographic assay of flunarizine, (*E*)-1-[bis(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)piperazine, in plasma of epileptic patients 444
Toyonaga, Y., see Ohmori, S. 15
Uralets, V.P., see Rodchenkov, G.M. 399
Van der Horst, F.A.L.
— and Holthuis, J.J.M.
Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 267
Van Schalkwijk, W.P., see Verkoelen, C.F. 41
Venkateswaran, K.S., see Ramachandran, P.K. 239
Verkoelen, C.F.
—, Romijn, J.C., Schroeder, F.H., Van Schalkwijk, W.P. and Splinter, T.A.W.
Quantitation of polyamines in cultured cells and tissue homogenates by reversed-phase high-performance liquid chromatography of their benzoyl derivatives 41
Vézina, D.
— and Bleau, G.
High-performance liquid chromatography of selenium in biological samples 385
Vijayaraghavan, R., see Ramachandran, P.K. 239
Watari, N., see Lee, F.W. 259
Weder, C.
—, Belsner, K., Lohkamp, R. and Höller, M.
Determination of prostaglandins D_2, E_2, $F_{2\alpha}$ and 6-ketoprostaglandin $F_{1\alpha}$ as well as thromboxane B_2 in rat brain by gas chromatography–mass spectrometry 345
Whelpton, R.
—, Jonas, G. and Buckley, D.G.
High-performance liquid chromatographic resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
Williams, P.E., see Brown, L.L. 370
Yonekawa, W.D., see Torchin, C.D. 444
Yoshimura, H., see Endoh, Y.S. 202
Zare, R.N., see Roach, M.C. 129
Zeman, I.
Misuse of the term "detergent" 452
Zschunke, M., see Brendel, E. 406

Subject Index

p-Acetamidobenzoic acid
Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 103
p-Acetamidohippuric acid
Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 103
Acetaminophen
Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 212
Acetaminophen glucuronide
Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 212
Acetaminophen sulphate
Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 212
Adenine nucleotides
Reversed-phase LC determination of cyclic adenosine 3′,5′-monophosphate in rat brain cortex 177
Adenosine phosphates
Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 249
n-Alkylamines
Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 267
Alphaprodine
Measurement of alphaprodine by selected-ion monitoring 392
p-Aminobenzoic acid
Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 103
p-Aminohippuric acid
Direct and simultaneous HPLC assay for the determination of *p*-aminobenzoic acid and its conjugates in human urine 103
δ-Aminolevulinic acid
Fluorimetric determination of urinary δ-aminolevulinic acid by HPLC and post-column derivatization 365
Androstenedione
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
Anilides
Determination of oxidation products of N-phenyllinoleamide: Spanish toxic oil syndrome studies 83
Atenolol
Simple extraction of atenolol from urine and its determination by HPLC 234
Benzimidazole
HPLC determination of formate as benzimidazole in biological samples 15
(*E*)-1-[(Bis(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)piperazine
HPLC assay of flunarizine, (*E*)-1-[bis(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)piperazine, in plasma of epileptic patients 444
Bisoprolol
Rapid and sensitive HPLC determination of bisoprolol in plasma and urine 431
β-Blockers
Qualitative GC and GC–MS screening for β-blockers in urine after solid-phase extraction using Extrelut-1 columns 194
Brobactam
Determination of 6β-bromopenicillanic acid (brobactam) in human serum and urine by HPLC 425
Bromofenofos
HPLC determination of bromofenofos and its metabolite in rat plasma 202
6β-Bromopenicillanic acid
Determination of 6β-bromopenicillanic acid (brobactam) in human serum and urine by HPLC 425
Cadralazine
Microbore LC determination of cadralazine and cephalexin in plasma with large-volume injection 335
Cephalexin
Microbore LC determination of cadralazine and cephalexin in plasma with large-volume injection 335
Cetyltrimethylammonium bromide
Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 267
Chiral stationary phases
Review. Direct separation of drug enan-

tiomers by HPLC with chiral stationary phases 1
Collagen α-chain polymers
Column LC separation of collagen α-chain polymers on sepiolite 162
Corticosteroids
Selective determination of urinary free cortisol by LC after solid-state extraction 25
Corticosteroids
Simplified method for the determination of plasma 18-hydroxycorticosterone 148
Corticosteroids
GC and MS study of synthetic corticosteroid metabolism: fluorometholone 399
Cortisol
Selective determination of urinary free cortisol by LC after solid-state extraction 25
Creatinine
Specific HPLC method for the determination of creatinine in rat plasma 358
Cyclic adenosine 3′,5′-monophosphate
Reversed-phase LC determination of cyclic adenosine 3′,5′-monophosphate in rat brain cortex 177
Dephosphate-bromofenofos
HPLC determination of bromofenofos and its metabolite in rat plasma 202
25-Desacetylrifampicin
Determination of rifampicin and its main metabolites in human plasma by HPLC 412
Desipramine
Simultaneous determination of imipramine, desipramine and their deuterium-labelled analogues in biological fluids by capillary GC–MS 93
Desmethylthioridazine
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
Detergent
Missue of the term "detergent" 452
2,3-Dihydro-6-[3-(2-hydroxymethyl)phenyl-2-propenyl]-5-benzofuranol
Determination of 2,3-dihydro-6-[3-(2-hydroxymethyl)phenyl-2-propenyl]-5-benzofuranol in plasma using LC with electrochemical detection 295
Dinucleotides
Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 249
Drugs
Review. Direct separation of drug enantiomers by HPLC with chiral stationary phases 1
Enantiomer separations
Review. Direct separation of drug enantiomers by HPLC with chiral stationary phases 1
Enantiomer separations
Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chiral β-cyclodextrin stationary phase 417
Enzymes
Microassay for GM_1 ganglioside β-galactosidase activity using HPLC 75
Enzymes
Hydrophobic chromatography: a one-step method for the purification of human liver microsomal epoxide hydrolase 169
Enzymes
Assay of γ-L-glutamylcyclotransferase activity in rat brain synaptosomes by HPLC 381
Epoxide hydrolase
Hydrophobic chromatography: a one-step method for the purification of human liver microsomal epoxide hydrolase 169
Estradiol
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
Estrone
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
Ethopropazine
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
Flunarizine
HPLC assay of flunarizine, (*E*)-1-[bis(4-fluorophenyl)methyl]-4-(3-phenyl-2-propenyl)piperazine, in plasma of epileptic patients 444
1-Fluoro-2,4-dinitrobenzene
Study of the derivatization of *n*-alkylamines with 1-fluoro-2,4-dinitrobenzene in the presence of aqueous cetyltrimethylammonium bromide micelles 267
Fluorometholone
GC and MS study of synthetic corticosteroid metabolism: fluorometholone 399
5-Fluorouracil
Determination of 5-fluorouracil in plasma and whole blood by ion-pair HPLC 121

Formate
HPLC determination of formate as benzimidazole in biological samples 15
3-Formyl-25-desacetylrifamycin SV
Determination of rifampicin and its main metabolites in human plasma by HPLC 412
3-Formylrifamycin SV
Determination of rifampicin and its main metabolites in human plasma by HPLC 412
Galabiosylceramide
TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga_2Cer 188
Galactosylceramide
TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga_2Cer 188
Glucosylceramide
TLC of neutral glycosphingolipids: an impoved simple method for the resolution of GlcCer, GalCer, LacCer and Ga_2Cer 188
γ-L-Glutamylcyclotransferase
Assay of γ-L-glutamylcyclotransferase activity in rat brain synaptosomes by HPLC 381
Glycocalicin
Human platelet glycocalicin purification by phenyl boronate affinity chromatography coupled to anion-exchange HPLC 376
Glycosphingolipids
TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga_2Cer 188
GM_1 ganglioside β-galactosidase
Microassay for GM_1 ganglioside β-galactosidase activity using HPLC 75
Guanine nucleotides
Reversed-phase LC determination of cyclic adenosine 3′,5′-monophosphate in rat brain cortex 177
Haemoglobin
GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 239
trans-6,6a,7,10,10a,11-Hexahydro-8,9-dimethyl-11-oxodibenz [*b,e*]oxepin-3-acetic acid
Enantiomeric analysis of a new anti-inflammatory agent in rat plasma using a chriral β-cyclodextrin stationary phase 417
High-mobility-group proteins
Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC: protein levels in rat tissues during postnatal development 65
Histone H1 subtypes
Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC; protein levels in rat tissues during postnatal development 65
Homovanillic acid
Measurement of homovanillic acid in small volumes of plasma using LC with electrochemical detection 141
Hormones
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
18-Hydroxycorticosterone
Simplified method for the determination of plasma 18-hydroxycorticosterone 148
7-Hydroxymethotrexate
Determination of methotrexate and its major metabolite, 7-hydroxymethotrexate, using capillary zone electrophoresis and laser-induced fluorescence detection 129
17α-Hydroxyprogesterone
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
20α-Hydroxyprogesterone
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
6-(1-Imidazolylmethyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid
Determination of the thromboxane synthetase inhibitor 6-(1-imidazolylmethyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (DP-1904) in human plasma and urine using solid-phase extraction and HPLC 111
Imipramine
Simultaneous determination of imipramine, desipramine and their deuterium-labelled analogues in biological fluids by capillary GC–MS 93
Immune complexes
Chromatographic and electrophoretic studies of immune complexes in non-A, non-B hepatitis 55
Immunoglobulins
Chromatographic and electrophoretic studies in immune complexes in non-A, non-B hepatitis 55

6-Ketoprostaglandin $F_{1\alpha}$
Determination of prostaglandins D_2, E_2, $F_{2\alpha}$ and 6-ketoprostaglandin $F_{1\alpha}$ as well as thromboxane B_2 in rat brain by GC–MS 345
Lactosylceramide
TLC of neutral glycosphingolipids: an improved simple method for the resolution of GlcCer, GalCer, LacCer and Ga_2Cer 188
Lanthionine ketimine
Specific method for the determination of 1,4-thiomorpholine-3,5-dicarboxylic acid and its unsaturated analogue lanthionine ketimine 183
Leucine
Rapid HPLC method to measure plasma leucine: importance in the study of leucine kinetics in vivo 370
LY 163502
HPLC assay of *trans*-(−)-5,5a,6,7,8,9,9a,10-octahydro-6-propylpyrimido[4,5-*g*]quinolin-2-amine dihydrochloride (LY 163502), a potent D-2 agonist, in plasma 438
Methotrexate
Determination of methotrexate and its major metabolite, 7-hydroxymethotrexate, using capillary zone electrophoresis and laser-induced fluorescence detection 129
Methotrimeprazine
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
3-Methoxy-4-hydroxyphenylglycol
Microassay of free 3-methoxy-4-hydroxyphenylglycol in plasma using HPLC with electrochemical detection 351
3-Methoxyparacetamol
Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 315
N-Methylcarbamylated haemoglobin
GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 239
R(−)10,11-Methylenedioxy-N-*n*-propylnoraporphine
HLPC separation and electrochemical or spectrophotometric determination of *R*(−)N-*n*-propylnorapomorphine and *R*(−)10,11-methylenedioxy-N-*n*-propylnorarphine in primate plasma 283
Methyl isocyanate
GC studies of the carbamylation of haemoglobin by methyl isocyanate in rats and rabbits 239
Nitrobenzylthioinosine
Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 249
Nitroglycerin
Simultaneous determination of nitroglycerin and its dinitrate metabolites by capillary GC with electron-capture detection 259
Nucleobases
Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 249
Nucleosides
Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 249
Nucleotides
Reversed-phase LC determination of cyclic adenosine 3′,5′-monophosphate in rat brain cortex 177
Nucleotides
Demonstration of the adenosine reservoirs with nitrobenzylthioinosine in liver and diaphragm by HPLC 249
trans-(−)-5,5a,6,7,8,9a,10,10-Octahydro-6-propylpyrimido [4,5-*g*]quinolin-2-amine dihydrochloride
HPLC assay of *trans*-(−)-5,5a,6,7,8,9,9a,10- octahydro-6-propylpyrimido[4,5-*g*]quinolin-2-amine dihydrochloride (LY 163502), a potent D-2 agonist, in plasma 438
Omeprazole
Improved procedure for quantitation of omeprazole and metabolites using reversed-phase HPLC 216
Omeprazole sulphide
Improved procedure for quantitation of omeprazole and metabolites using reversed-phase HPLC 216
Omeprazole sulphone
Improved procedure for quantitation of omeprazole and metabolites using reversed-phase HPLC 216
Oxprenolol
Determination of oxprenolol in plasma by column LC 229
Paracetamol

Quantitation of acetaminophen and its metabolites in rat plasma after a toxic dose 212
Paracetamol
Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 315
Phenobarbital
Simultaneous determination of phenytoin and phenobarbital in human serum by electrochemical immunoassay combined with HPLC 449
N-Phenyllinoleamide
Determination of oxidation products of N-phenyllinoleamide: Spanish toxic oil syndrome studies 83
Phenytoin
Simultaneous determination of phenytoin and phenobarbital in human serum by electrochemical immunoassay combined with HPLC 449
Polyamines
Quantitation of polyamines in cultured cells and tissue homogenates by reversed-phase HPLC of their benzoyl derivatives 41
Prenylamine
Simultaneous determination of *R*- and *S*-prenylamine in plasma and urine by reversed-phase HPLC 305
Progesterone
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
Promethazine
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
R(−)N-*n*-Propylnorapomorphine
HPLC separation and electrochemical or spectrophotometric determination of *R*(−)N-*n*-propylnorapomorphine and *R*(−)10,11-methylenedioxy-N-*n*-propylnoraporphine in primate plasma 283
Prostaglandins
Determination of prostaglandins D_2, E_2, $F_{2\alpha}$ and 6-ketoprostaglandin $F_{1\alpha}$ as well as thromboxane B_2 in rat brain by GC–MS 345
Proteins
Separation and quantification of histone H1 subtypes and high-mobility-group proteins by reversed-phase LC: protein levels in rat tissues during postnatal development 65
Pseudoephedrine
Sensitive HPLC determination of pseudoephedrine in plasma and urine 406
Rifampicin
Determination of rifampicin and its main metabolites in human plasma by HPLC 412
Selenium
HPLC of selenium in biological samples 385
Steroids
Selective determination of urinary free cortisol by LC after solid-state extraction 25
Steroids
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
Steroids
Simplified method for the determination of plasma 18-hydroxycorticosterone 148
Steroids
GC and MS study of synthetic corticosteroid metabolism: fluorometholone 399
Testosterone
Automated direct assay system for the measurement of sex steroid hormones in serum using HPLC 33
3-Thiomethylparacetamol
Complete separation of urinary metabolites of paracetamol and substituted paracetamols by reversed-phase ion-pair HPLC 315
1,4-Thiomorpholine-3,5-dicarboxylic acid
Specific method for the determination of 1,4-thiomorpholine-3,5-dicarboxylic acid and its unsaturated analogue lanthionine ketimine 183
Thioridazine
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
Thromboxane B_2
Determination of prostaglandins D_2, E_2, $F_{2\alpha}$ and 6-ketoprostaglandin $F_{1\alpha}$ as well as thromboxane B_2 in rat brain by GC–MS 345
Thromboxane synthetase inhibitor
Determination of the thromboxane synthetase inhibitor 6-(1-imidazolylmethyl)-5,6,7,8-tetrahydronaphthalene-2-carboxylic acid (DP-1904) in human plasma and urine using solid-phase extraction and HPLC 111
Trimeprazine

HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
Trimipramine
HPLC resolution of the enantiomers of thioridazine, its metabolites and related compounds 223
Tyrosines
Analysis of the isomeric tyrosines in mammalian and avian systems using HPLC with fluorescence detection 155
Xylazine
Determination of xylazine in blood components using HPLC 207